

# Real Analysis 4-5: The Integral Test for Series

Brad Baxter  
Birkbeck College, University of London

June 15, 2023

You can download these slides and the lecture videos from my office server

<http://econ109.econ.bbk.ac.uk/brad/analysis/>

**Recommended books:** Lara Alcock (2014), “How to Think about Analysis”, Oxford University Press.

J. C. Burkill (1978), “A First Course in Mathematical Analysis”, Cambridge University Press.

Riemann integration is studied in depth in Lecture 7.

**BUT** there is a simple Integral Test that relates series to integral of functions for which you only need to know that decreasing and increasing functions are Riemann integrable and have the following simple property:

### Theorem

If  $f_1(x) \geq f_2(x)$  for  $x \in [a, b]$ , then

$$\int_a^b f_1(x) dx \geq \int_a^b f_2(x) dx.$$

This lecture will enable you to complete Q7 on HW3.

## Theorem (Integral Test for series convergence)

Let  $f: [1, \infty) \rightarrow (0, \infty)$  be a **positive decreasing** function. Then

①

$$\sum_{n=1}^{\infty} f(n) \quad \text{and} \quad \int_1^{\infty} f(x) dx$$

either both diverge or both converge.

② Let

$$a_n = \sum_{k=1}^n f(k) - \int_1^n f(x) dx, \quad n \in \mathbb{N}.$$

Then  $(a_n)$  is a decreasing sequence and  $0 \leq a_n \leq f(1)$ , so it's convergent with limit in  $[0, f(1)]$ .

## Proof.

- ① If  $k - 1 \leq x \leq k$ , then  $f(k - 1) \geq f(x) \geq f(k)$ . Integrating over  $(k - 1, k)$ ,

$$f(k - 1) \geq \int_{k-1}^k f(x) dx \geq f(k).$$

Adding these inequalities

$$\sum_{k=1}^{n-1} f(k) \geq \int_1^n f(x) dx \geq \sum_{k=2}^n f(k). \quad (*)$$

Hence  $\int_1^\infty f$  and  $\sum_1^\infty f(n)$  both converge or both diverge.

- ② We have

$$a_n - a_{n-1} = f(n) - \int_{n-1}^n f(x) dx \leq 0$$

and  $0 \leq a_n \leq f(1)$  by  $(*)$  (**exercise**).

## Example ( $\sum n^{-3/2}$ convergent)

The series

$$\sum_{n=1}^{\infty} \frac{1}{n^{3/2}}$$

is convergent if and only if the infinite integral

$$\int_1^{\infty} x^{-3/2} dx$$

exists. But

$$\int_1^A x^{-3/2} dx = \left[ -2x^{-1/2} \right]_1^A = 2 - \frac{2}{A^{1/2}} \rightarrow 2,$$

as  $A \rightarrow \infty$ . Hence

$$\sum_{n=1}^{\infty} \frac{1}{n^{3/2}}$$

is convergent.

### Example (Harmonic series $\sum 1/n$ is divergent)

Let  $f(x) = 1/x$ . Here's the Integral Test in detail (draw a picture!):

$$S_N = \sum_{n=1}^N \frac{1}{n} \geq \int_1^{N+1} \frac{1}{x} dx = \ln(N+1).$$

Since  $\ln(N+1) \rightarrow \infty$  as  $N \rightarrow \infty$ , the partial sums ( $S_N$ ) are unbounded. Hence the harmonic series is divergent.

**Logarithms:** Here we are using

$$\ln y = \int_1^y \frac{1}{x} dx, \quad \text{for } y > 0,$$

which we shall discuss in Lecture 7.

Example ( $\sum 1/n^c$  is divergent for  $0 < c < 1$ )

Here's the Integral Test in detail again (draw a picture!):

$$S_N = \sum_{k=1}^N \frac{1}{n^c} \geq \int_1^{N+1} x^{-c} dx = \left[ \frac{x^{1-c}}{1-c} \right]_1^{N+1} = \frac{(N+1)^{1-c} - 1}{1-c}.$$

But, if  $c < 1$ , then  $1 - c > 0$  and  $(N + 1)^{1-c} \rightarrow \infty$  as  $N \rightarrow \infty$ . Thus the partial sums ( $S_N$ ) are unbounded and the series is divergent.



Example ( $\sum 1/n^c$  is convergent for  $c > 1$ )

Here's the Integral Test in detail again (draw a picture!):

$$\sum_{k=2}^N \frac{1}{n^c} \leq \int_1^N x^{-c} dx = \left[ \frac{x^{1-c}}{1-c} \right]_1^N = \frac{N^{1-c} - 1}{1-c}.$$

But, if  $c > 1$ , then  $1 - c < 0$  and  $N^{1-c} \rightarrow 0$  as  $N \rightarrow \infty$ . Thus the partial sums ( $S_N$ ) are bounded and the series is convergent.

## Theorem (Euler's constant)

Let

$$a_n = 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} - \log n.$$

Then  $a_n \rightarrow \gamma$ , where  $0 < \gamma < 1$ .

Proof.

Apply the Integral Test to  $f(x) = 1/x$ . □